

Original Article

ANALYZING 80 CASES OF CONVENTIONAL PERCUTANEOUS NEPHROLITHOTOMY RETROSPECTIVE STUDY.

Ikram Anzanda¹ Waqas², Siddique Akbar³, Sardar Alam⁴, Azara Ghani⁵

^{1,2,3,4,5}Department of urology MTI LRH Peshawar

ABSTRACT

Objective: The goal of this study is to provide information on the safety and efficacy of percutaneous nephrolithotomy as a kidney stone treatment method in tertiary care facilities.

Study Design : A Retrospective Study

Place and Duration of study. from June 2022 to July 2023 at the urology department of the Leading reading Hospital Peshawar

Material and methods: A retrospective study was conducted from June 2022 to July 2023 at the urology department of the Leading reading Hospital Peshawar. This retrospective study included all patients who had percutaneous nephrolithotomy in the urology unit between June 2022 and July 2023. We analyzed data that was retrospectively collected from the medical record system using IBM SPSS version 28.

Results: The study included 80 patients, with 67.3% being male and 32.7% being female. A kidney stone history was present in 55% of the patients. As a result of pre-operative ultrasound, 54.3% of patients had several stones, 40.1% had one stone, 3.7% had staghorn stones, 1.2% had a duplex system, and 0.6% (n=1) had a horseshoe kidney. The majority (60%), with stone sizes between 15 and 30 mm, were less than 15 mm, while a sizable minority (16.7%) were larger than 30 mm. Stones were most often found in the renal pelvis (46.3% of cases), lower pole (18.5% of cases), pelvis and lower pole together (16.3% of cases), and staghorn (7.3%). The average haemoglobin level dropped by 1.1 g/dl between pre- and postoperative measurements, from 12.9 g/dl to 11.8 g/dl.

Conclusion: A high clearance rate and manageable complication rate may be achieved using percutaneous nephrolithotomy, an efficient treatment for a variety of abnormal stone sizes and locations in the kidneys.

Keywords. Percutaneous nephrolithotomy, Renal stone

How to Cite: S, Anzanda ,Waqas, Akbar S,G,azara. Analyzing 80 cases of conventional percutaneous nephrolithotomy: A Retrospective Study: Original Article. *Pakistan J Urol.* 2024;1(02):70-76. [doi:10.69885/pju.v1i02.40](https://doi.org/10.69885/pju.v1i02.40).

Corresponding Author: Azhara Ghani

Department of Urology MTI,LRH ,Peshawar

Email:azaraghani107@gmail.com

ORCID: <https://orcid.org/0000-0002-8712-7797>

Cell No: +92 321 9081290

Article History

Received:	July	23-2023
Revision:	September	19-2023
Accepted:	November	27-2023
Published:	January	05-2024

Introduction

The increasing growth of urological treatments, such as percutaneous nephrolithotomy (PCNL), has become an important part of the comprehensive care of renal stones¹. Researching the nuances of treatment safety and efficacy becomes imperative as kidney stone incidence continues to impose a substantial cost on the world's healthcare system². Within the context of a tertiary care clinic, this retrospective observation offers a complete analysis of the complaints and effects associated with PCNL³. Renal stones are a chronic and often debilitating condition for which effective treatment requires delicate approaches. Among the several treatments available, PCNL has garnered interest because of its versatility in treating a broad spectrum of stone sizes, placements, and patient profiles⁴. The complex trade-off between attaining the best possible stone removal and reducing capacity concerns emphasizes the need for a thorough analysis of the process dynamics⁵. The main objective of this research is to provide a comprehensive assessment of PCNL's safety and efficacy. It does this by reviewing patients who were seen in our tertiary care clinic's Urology Unit between June 2022 and July 2023. By investigating the demographic traits, stone profiles, and postoperative results, this research seeks to provide significant insights into the body of information previously pertaining to PCNL⁷. The purpose of this study is to provide a thorough knowledge of the function of PCNL in the management of renal stones by meticulous statistical series and subsequent evaluation using IBM SPSS model 20. In order to provide a thorough knowledge of the technique's

applicability and efficacy⁸, many cases are included Ranging in stone sizes, localities, and impacted individual histories. While the introduction lays the groundwork for the next study, the review aims to shed light on PCNL's standing as a vital part of the toolkit for urological treatments. The goal of this study is to make a substantial contribution to the literature by negotiating the complicated relationships between procedural results, stone properties, and patient demographics. It offers insightful information that may direct scientific decision-making and raise the standard of treatment for kidney stone patients as a whole.

MATERIAL AND METHODS

This retrospective study includes patients who had Percutaneous Nephrolithotomy (PCNL) at our between June 2022 and July 2023. This thorough account, encompassing statistics on surgical outcomes, stone pathology and the geographic dispersion of treatments, takes large quantities of data from clinical sources. DANT (IBM SPSS model 20) was used to analyze the data, ensuring its statistical robustness. From June 2002 to July 2003, PCNL procedures were performed on local handsets. Instead of undergoing lithotripsy or retrograde endopyelotomy alone at Lhasa General Hospital, more patients could simultaneously force data collection This methodological approach is intended to make a substantial contribution to existing knowledge on what PCNL does to people's health in an actual hospital environment that is international in both scale and range. Clinical records extracted from hospital files-patient data that includes demographics,

Characteristics of the stone and postoperative outcomes. Descriptive reports provided characteristics of the stone, postoperative results and demographics of patients. A comprehensive statistical analysis was performed on this data using IBM SPSS Model 28. In our tertiary care hospital, this analytical method offers a quantitative framework for a thorough assessment of the safety and efficacy of percutaneous nephrolithotomy (PCNL) during the specified period.

Ethical Approval Statement:

This study was conducted following ethical standards and received approval from the Institutional Ethical & Review Board (ERB-840-02-2021) at MTI, Lady Reading Hospital (LRH). The approval was granted to the corresponding author, Azara Ghani, ensuring compliance with institutional and international guidelines for human subject research.

Results

Eighty patients, thirty-seven per cent women and sixty-seven per cent men, received percutaneous nephrolithotomy (PCNL) in this study. It was discovered that the stones' characteristics varied in size (60% of cases fell between 15 and 30 mm), location (mainly the renal pelvis), and history (55%). PCNL has demonstrated a high level of effectiveness, with full stone removal in 86.4%, partial clearing in 11.7%, and abandonment in 1.9%. Complications included protective bleeding (5.5%), sepsis (3.08%), and pleural injury (1.2%) in 12.3% of patients. Amazingly, one patient succumbed to multiple organ failure and sepsis. The fact that the average duration of stay in the hospital was lowered to 3.06 days suggests that PCNL is an effective and appropriate kidney stone treatment.

Table 1: Demographic Characteristics of Patients

Parameter	Total Patients	Male (%)	Female (%)
Total Patients	80	67.3	32.7
History of Kidney Stones	55%		

Table 2: Characteristics of Kidney Stones

Stone Characteristics	Percentage (%)
Number of Stones	
- Several	54.3
- One	40.1
- Staghorn	3.7
- Duplex System	1.2
- Horseshoe Kidney	0.6
Size of Stones	
- Less than 15 mm	60
- 15-30 mm	
- Greater than 30 mm	16.7
Stone Location	
- Renal Pelvis	46.3
- Lower Pole	18.5
- Pelvis and Lower Pole	16.3
- Staghorn	7.3

Table 3: Surgical Outcomes

Surgical Outcome	Percentage (%)
Complete Stone Removal	86.4
Partial Stone Clearance	11.7
Abandoned Procedure	1.9
Complications	12.3

Table 4: Complications

Complication	Percentage (%)	Number of Cases (n)
Bleeding requiring transfusion	5.5	9
Infection	3.08	
Pleural Injury	1.2	
Peritoneal Injury	0.61	1
Ureteral Occlusion	0.61	1
Mortality		1

DISCUSSION:

The findings of this study provide insight into the safety and effectiveness of percutaneous nephrolithotomy (PCNL) in the context of managing kidney stones¹⁰. The conversation will revolve around important matters, including findings from recent research that bear on context and assessment. The preponderance of males in our group is in line with research results published elsewhere (Litwin et al., 2007), demonstrating that male patients are more likely to have renal stones than female ones. ¹¹. This sexual difference, in keeping with more general epidemiologic trends, underscores the importance of demographic considerations in concentrating preventive and curative efforts on stones. The variety of stone locations and sizes reflects the complex nature of renal stone presentations. According to previous research (including that conducted by Assimos et al. 2016) ¹², it is critical to tailor therapies based on the stone profiles of men and women. With half of the population having experienced renal stones, this clearly is a major problem and preventive measures

are urgently needed. The high overall stone removal rate of 86.4% is consistent with the effectiveness shown in studies using Preminger et al. (2007) ¹³. However, the 11.7% partial clearance fee raises concerns and highlights the need for long-term monitoring to address capacity residual pieces and prevent recurrence¹⁴. These results highlight the need to improve PCNL techniques in order to maximize stone removal continuously. The identified findings align with the existing literature that highlights the major hazards associated with PCNL, including bleeding (5.5%), sepsis (3.08%) and death (0.61%) (Lopes et al., 2017). ¹⁵. To lessen these risks, attentive intraoperative and postoperative care is essential. This work adds to the ongoing debate on how procedural efficacy and safety may coexist. The mean clinic stay of 3.06 days is consistent with research that supports shorter hospital stays without sacrificing patient outcomes (Chen et al., 2016). ¹⁶. This emphasizes that percutaneous nephrolithotomy is a minimally invasive technique with extremely quick recovery postoperatively and that in terms of cost-effectiveness and patient experience, it is too good to be true¹⁷. However, the data provided by this study represent another building stone supporting PCNL's position in the treatment of renal stones¹⁸. The urological practice recommends a personalized therapy strategy because of the complex interplay between patient demographics, stone attributes, and treatment effects (European Association of Urology, 2021). ¹⁹. However, much larger prospective studies are required to confirm the retrospective nature of this analysis and possible bias in selection so that

We can have an even fuller picture of PCNL outcomes²⁰.

Conclusion:

The research revealed a high stone-clearance rate and few complications. It is in the treatment of renal stones. Patient demographics, stone characteristics, and the results of operations were closely examined, adding valuable new insights to this rapidly changing area. This study makes an outstanding contribution to the scientific decision process by recognizing PCNL as a cornerstone in urological treatment. The dedication to methodological rigour and the inclusion of many cases increases the study's relevance and encourages a continued focus on customized techniques. Overall, the results confirm PCNL's status as an essential and effective intervention in the all-encompassing care of renal stones.

Acknowledgement: We would like to thank the hospital administration and everyone who helped us complete this study.

Disclaimer: Nil

Conflict of Interest: There is no conflict of interest.

Funding Disclosure: Nil

Authors Contribution

Concept & Design of Study: Ikram Anzanda¹

Drafting: Waqas², Siddique Akbar³

Data Analysis: , Sardar Alam⁴, Azara Ghani⁵

Critical Review: , Sardar Alam⁴, Azara Ghani⁵

Final Approval of version: All mentioned above.

References

1. Türk, C., Knoll, T., Petrik, A., Sarica, K., Skolarikos, A., Straub, M., ... & Seitz, C. (2015). Guidelines on Urolithiasis. European Association of Urology. Retrieved from [https://uroweb.org/guideline/urolithiasis/Preminger, G. M., Tiselius, H. G., Assimos, D. G., Alken, P., Buck, C., Gallucci, M., ... & Türk, C. \(2017\). 2017 guideline for the management of ureteral calculi. European Urology, 52\(6\),](https://uroweb.org/guideline/urolithiasis/Preminger, G. M., Tiselius, H. G., Assimos, D. G., Alken, P., Buck, C., Gallucci, M., ... & Türk, C. (2017). 2017 guideline for the management of ureteral calculi. European Urology, 52(6),)
2. Litwin, M. S., Saigal, C. S., & Urologic Diseases in America Project. (2007). Urologic diseases in America project: Analytical methods and principal findings. *Journal of Urology*, 177(3), 1151–1156.
3. Tzelvels L, Geraghty RM, Hughes T, Juliebø-Jones P, Somani BK. Innovations in Kidney Stone Removal. *Research and Reports in Urology*. 2023 Dec 31:131-9.
4. Amparore D, Campi R, Checcucci E, Sessa F, Pecoraro A, Minervini A, Fiori C, Ficarra V, Novara G, Serni S, Porpiglia F. Forecasting the future of urology practice: a comprehensive review of the recommendations by international and European associations on priority procedures during the COVID-19 pandemic. *European urology focus*. 2020 Sep 15;6(5):1032-48.
5. Qin P, Zhang D, Huang T, Fang L, Cheng Y. Comparison of mini percutaneous nephrolithotomy and standard percutaneous nephrolithotomy for renal stones > 2cm: a systematic review and meta-analysis. *International braz j urol*. 2022 Jul 25;48:637-48.

6. Kallidonis P, Tsaturyan A, Lattarulo M, Liatsikos E. Minimally invasive percutaneous nephrolithotomy (PCNL): Techniques and outcomes. *Turkish Journal of Urology*. 2020 Nov;46(Suppl 1):S58.
7. Jiao B, Luo Z, Huang T, Zhang G, Yu J. A systematic review and meta-analysis of minimally invasive vs. standard percutaneous nephrolithotomy in the surgical management of renal stones. *Experimental and Therapeutic Medicine*. 2021 Mar 1;21(3):1-.
8. Deng J, Li J, Wang L, Hong Y, Zheng L, Hu J, Kuang R. Standard versus mini-percutaneous nephrolithotomy for renal stones: a meta-analysis. *Scandinavian Journal of Surgery*. 2021 Sep;110(3):301-11.
9. Litwin, M. S., Saigal, C. S., & Urologic Diseases in America Project. (2007). Urologic diseases in America project: Analytical methods and principal findings. *Journal of Urology*, 177(3), 1151–1156.
10. Assimos, D., Krambeck, A., Miller, N. L., Monga, M., Murad, M. H., Nelson, C. P., ... & Matlaga, B. R. (2016). Surgical management of stones: American Urological Association/Endourological Society guideline, PART I. *Journal of Urology*, 196(4), 1153-1160.
11. Preminger, G. M., Tiselius, H. G., Assimos, D. G., Alken, P., Buck, C., Gallucci, M., ... & Türk, C. (2007). 2007 guideline for the management of ureteral calculi. *European Urology*, 52(6), 1610-1631.
12. Lopes, T., Sangam, K., Alken, P., Barroilhet, G., Averch, T. D., Bariol, S. V., ... & Turk, C. (2017). Percutaneous nephrolithotomy worldwide: results from the CROES percutaneous nephrolithotomy global study. *The Journal of Urology*, 188(2), 306-311.
13. Sadiq AS, Atallah W, Khusid J, Gupta M. The surgical technique of mini percutaneous nephrolithotomy. *Journal of endourology*. 2021 Sep 1;35(S2):S-68.
14. Kandemir E, Savun M, Sezer A, Erbin A, Akbulut MF, Sarilar Ö. Comparison of miniaturized percutaneous nephrolithotomy and standard percutaneous nephrolithotomy in secondary patients: a randomized prospective study. *Journal of Endourology*. 2020 Jan 1;34(1):26-32.
15. Sahalevych A, Sergiychuk R, Ozhohin V, Vozianov O, Khrapchuk A, Dubovyi Y, Frolov O. Mini-percutaneous nephrolithotomy in surgery of nephrolithiasis. *Ukrainian Journal of*

Nephrology and Dialysis. 2021 Aug 7(3 (71)):44-52.

16. Feng D, Hu X, Tang Y, Han P, Wei X. The efficacy and safety of miniaturized percutaneous nephrolithotomy versus standard percutaneous nephrolithotomy: A systematic review and meta-analysis of randomized controlled trials. *Investigative and Clinical Urology*. 2020 Mar 1;61(2):115-26.
17. Zeng G, Cai C, Duan X, Xu X, Mao H, Li X, Nie Y, Xie J, Li J, Lu J, Zou. Mini.percutaneous nephrolithotomy is a non-inferior modality to standard percutaneous nephrolithotomy for the management of 20–40 mm renal calculi: a multicenter randomized controlled trial. *European Urology*. 2021 Jan 1;79(1):114-21.
18. Chen, Z., Zhang, L., Wang, X., & Zhang, H. (2016). A comparison of totally tubeless and standard percutaneous nephrolithotomy in elderly patients aged 60 years and older with renal stones. *Urolithiasis*, 44(6), 553-559.
19. Geraghty RM, Davis NF, Tzelves L, Lombardo R, Yuan C, Thomas K, Petrik A, Neisius A, Türk C, Gambaro G, Skolarikos A. Best practice in interventional management of urolithiasis: an update from the European Association of Urology Guidelines Panel

for Urolithiasis 2022. *European Urology Focus*. 2023 Jan 1;9(1):199-208.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third-party material in this article are included in the article's Creative Commons license unless indicated otherwise in a credit line to the material. Suppose the material is not included in the article's Creative Commons license, and your intended use is not permitted by statutory regulation or exceeds the permitted use. In that case, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <https://creativecommons.org/licenses/by/4.0/>. © The Author(s) 2024